

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/372344158>

An Oriental Pratincole *Glareola maldivarum* egg in a Black-winged Stilt *Himantopus himantopus* nest

Article in *Indian BIRDS* · July 2023

CITATIONS

0

READS

17

2 authors:



Shalini Jain

National Cheng Kung University

3 PUBLICATIONS 0 CITATIONS

SEE PROFILE



Mayank Shukla

National Centre for Biological Sciences, Bangalore

3 PUBLICATIONS 1 CITATION

SEE PROFILE

Varu, S.N., 1988. Some information on the shorebirds of Kutch. *Newsletter for Birdwatchers* 28: 13.

Varu, M. & Varu, S., 2018. Tagged Great Knot from Jamnagar. *Flamingo*. XVI-4. Pp. 12-13

Ward, S., 2012. *Great Knot Calidris tenuirostris. Threatened Species of the Northern Territory*. Published by Northern Territory Government. Pp. 1-3.

– **Yashodhan Bhatia, Ashish Pankhania, Mehul Bhadania & Dishant Parasharya**

Yashodhan Bhatia, Nirmal Foundation frd, Jamnagar. 361008.

E-mail: jamnagarbirds@gmail.com (Corresponding Author)

Ashish Pankhania, 101/ Madhav Kunj apt., Vasa Vira society, street # 3, near Under-bridge, Jamnagar 361006. E-mail: ashish.mr.birds@gmail.com

Dr. Mehul Bhadania, 53 / A, AmbaVijay Society, Opp Mahavir C, Jamnagar 361008.

E-mail: mehulbhadania@gmail.com

Dr. Dishant Parasharya Scientist B, Bombay Natural History Society, Mumbai 400 001.

E-mail: dparasharya@gmail.com

An Oriental Pratincole *Glareola maldivarum* egg in a Black-winged Stilt *Himantopus himantopus* nest

On 6 June 2022, at 1006 h, while returning after bird watching at Bhigwan, Pune (18.28°N, 74.77°E), we came across a nest with three Black-winged Stilt *Himantopus himantopus* and one Oriental Pratincole *Glareola maldivarum* egg [34]. The nest was located in an unused agricultural field. The nest was a shallow mud scrape made up of shell fragments, twigs, and agricultural stubble. The Black-winged Stilt eggs were cream-coloured with black and brown spots, whereas the Oriental Pratincole egg was relatively smaller in size with black-brown blotches on a pale brown base.



Both: Mayank Shukla

34. Black-winged Stilt nest with an egg of Oriental Pratincole.

The extra egg presumed as the Oriental Pratincole egg based on our observations of several other Oriental Pratincoles incubating in the same agricultural field, and the egg was visually compared with other Oriental Pratincole eggs. Photographs were taken from a safe distance without approaching very close to the nest. After photographing the nest, we moved outside the field carefully to a distant point to observe the nest using binoculars. The observations were made from a distance of about 15 m. We lied down on the ground after scanning the sky and nearby area to ensure that there are no predators and then observed the nest. After a few minutes, a pair of Black-winged Stilt approached the nest and one of the birds started to incubate while the other one flew away [35]. We observed the nest for about one hour during which the eggs were continuously incubated by the parent bird

[35]. We could only observe the same nest the next day for about 15 min and found no change in egg composition and number. The Black-winged Stilt pair was found to incubate the nest.



35. Parent Black-winged Stilt incubating.

The breeding ecology of Oriental Pratincole has not been studied extensively (Maclean & Kirwan 2020), however further emphasis on this could potentially reveal if such instances are just mistakes of egg laying by Oriental Pratincole or possibly a case of brood parasitism.

We are thankful to Tamás Székely for his support.

References:

Maclean, G.L., & Kirwan, G.M., 2020. Oriental Pratincole (*Glareola maldivarum*), version 1.0. In *Birds of the World* (J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, and E. de Juana, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. Webpage URL: <https://doi.org/10.2173/bow.oripra.01>. [Accessed on 20 October 2022.]

– **Mayank Shukla & Shalini Jain**

Mayank Shukla, 1407, Victory-B, Nirmal Lifestyle, Kalyan (W), Thane 421102, Maharashtra, India. Email: mayankshuklams@gmail.com [Corresponding author]
Shalini Jain, 9/85, Chhippaity, Aligarh 200201, Uttar Pradesh, India.
Email: jshaliniwls@gmail.com

High altitude records of Orange-headed Thrush *Geokichla citrina* and Asian Koel *Eudynamis scolopaceus* in the Nepal Himalaya

We report sightings of Orange-headed Thrush *Geokichla citrina* and Asian Koel *Eudynamis scolopaceus* from altitudes previously unreported for them, from ward number 4 of Gharapjhong Rural Municipality, Mustang District, Gandaki Province, Nepal, which lies in the Annapurna Conservation Area.

In Nepal, the Orange-headed Thrush is a locally common and mainly a summer visitor, with some birds remaining all year as widespread partial migrants (Inskipp et al. 2016). This bird inhabits wet ravines and understorey of the moist deciduous forests (Inskipp & Inskipp 1991), evergreen forest, bamboo thickets, and plantations in tropical and subtropical zones (Collar & Juana 2020). The maximum elevations documented for this species are 1,830 m (Ali & Ripley 1971; Kazmeirczak 2000 ; Rasmussen & Anderton 2012; Inskipp et al. 2016; Grimmer et al. 2016), and 2,300 m (BirdLife International 2022a).

On Saturday, 22 May 2021, one Orange-headed Thrush entered a house in Jomsom village (28.78°N, 83.73°E) and got